

51) International Patent Classification 5:		11) International Publication Number: WO	93/1177
A61K 31/57 // (A61K 31/57 A61K 31:165)	A1	43) International Publication Date: 24 June 199	3 (24.06.93
21) International Application Number: PCT/E.		DE, DK, ES, FI, GB, HU, JP, KP, KR, LK	, LU, MO
22) International Filing Date: 7 December 1992	. (07.12.	European patent (AT, BE, CH, DE, DK, E GR, IE, IT, LU, MC, NL, PT, SE), OAPI BJ, CF, CG, CI, CM, GA, GN, ML, MR, SN	S, FR, GI patent (Bl
30) Priority data: 91311761.0 18 December 1991 (18.1	2.91)	Published	, 12, 10,
71) Applicant: AKTIEBOLAGET ASTRA [SE/SE] Södertälje (SE).		With international search report. Before the expiration of the time limit for an claims and to be republished in the event of t.	nending th he receipt o
72) Inventors: CARLING, Christer, Carl, Gustav; 8, S-240 10 Dalby (SE). TROFAST, Jan, Wil penkroken 34, S-226 47 Lund (SE).	Backväg liam; \	amendments.	
74) Agents: HJERTMAN, Ivan et al.; AB Astra, l partment, S-151 85 Södertälje (SE).	Patent I	-	
•			
54) Title: NEW COMBINATION OF FORMOTE	ROL A	D BUDESONIDE	
57) Abstract			
Effective amounts of formoterol (and/or a phy n combination for simultaneous, sequential or separate	vsiologic rate adn	ly acceptable salt and/or solvate thereof) and budesoni nistration by inhalation in the treatment of respiratory	de are use disorder.
· ·			
		·	

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

FOR THE PURPOSES OF INFORMATION ONLY

AT AU BB BE BF BG BJ BR CA CF CG CH CI CM CS CZ DE DK ES	Austria Australia Barbados Belgium Burkina Faso Bulgaria Benin Brazil Canada Central African Republic Congo Switzerland Côte d'Ivoire Cameroon Czechoslovakia Czech Republic Germany Denmark Spain Finland	FR GA GB GN GR HU IE IT JP KP KR KZ LJ LK LU MC MG ML MN	France Gabon United Kingdom Guinea Greece Hungary Ireland Italy Japan Democratic People's Republic of Korea Republic of Korea Kazaklistan Liechtenstein Sri Lanka Luxembourg Monaco Madagascar Mali Mongolia	MR MW NL NO NZ PL PT RO RU SD SE SK SN TD TG UA US VN	Mauritania Malawi Netherlands Norway New Zealand Poland Portugal Romania Russian Federation Sudan Sweden Slovak Republic Senegal Soviet Union Chad Togo Ukraine United States of America Viet Nam
--	--	--	--	---	---

New combination of formoterol and budesonide.

5

10

15

Field of the invention

This invention relates to improvements in the treatment of mild as well as severe asthma and other respiratory disorders. More particularly, it relates to the use of a bronchodilator in combination with a steroidal anti-inflammatory drug for the treatment of respiratory disorders such as asthma, and to pharmaceutical compositions containing the two active ingredients. It emphasizes the use of a long-acting bronchodilator which provides rapid relief of symptoms.

Background of the invention

There have recently been significant advances in our 20 understanding of asthma. Despite many advances, both in awareness of the disease by doctors and patients alike, coupled with the introdction of very powerful and effective anti-asthma drugs, asthma remains a poorly 25 understood and often poorly treated disease. Previously, contraction of airway smooth muscles has been regarded as the most important feature of asthma. Recently there has been a marked change in the way asthma is managed, stemming from the fact that asthma is recognized as a chronic inflammatory disease. Uncontrolled airway 30 inflammation may lead to mucosal damage and structural changes giving irrversible narrowing of the airways and fibrosis of the lung tissue. Therapy should therefore be aimed at controlling symptoms so that normal life is 35 possible and at the same time provide basis for treating

the underlying inflammation.

The most common cause for poor control of asthma is poor compliance with the long-term management of chronic asthma, particularly with prophylatic treatments, such as inhaled steroids, which do not give immediate symptom relief. Patients will readily take β_2 -agonist inhalers, since these provide rapid relief of symptoms, but often do not take prophylactic therapy, such as inhaled steroids, regularly because there is no immediate symptomatic benefit. They also counteract down regulation of β_2 -adrenoceptor agonists.

methoxyphenyl)-1-methylethyl]amino]ethyl]phenyl] 15 formamide), is an adrenoceptor agonist which selectively stimulates β_2 -receptors, thus producing relaxation of bronchial smooth muscle, inhibition of the release of endogenous spasmogens, inhibition of oedema caused by endogenous mediators, and increased mucociliary 20 clearence. Inhaled formoterol fumarate acts rapidly, usually within minutes which gives the patient immediate confirmation that he has taken an adequat dose and thereby avoiding overdosing of both β -agonist and steroid. Inhaled formoterol also exerts a prolonged 25 bronchodilation, which in clinical trials has been demonstrated as up to 12 hours.

Budesonide, (16,17-butylidenebis(oxy)-11,21
dihydroxypregna-1,4-diene-3,20-dione), may be given in a high inhaled dose (up to 2 mg daily) with very low systemic effects, possibly because of its rapid metabolism. The high rapid systemic elimination of budesonide is due to extensive and rapid hepatic

metabolism. Long term clinical studies have shown that inhaled budesonide is a pharmacologically safe drug. High doses of inhaled budesonide are highly effective and well

3

tolerated when used in oral steroid replacement therapy. Budesonide represents a logical safe and effective therapy for long term control of asthma.

- 5 The inhaled route of administration enables the dose to be delivered directly to the airways. By this type of administration, it is possible to give a small dose and thereby minimizing unwanted side-effects. The drawbacks of the currently available bronchodilators are their relatively short duration of action. By using a compound with long duration e.g. formoterol it would be possible to avoid the nocturnal asthma, which so often causes considerable anxiety and debility to the patients. Formoterol gives less nocturnal waking than the commonly used short-acting agonists like salbutamol, terbutaline and the like. Formoterol has been registered for oral
- Pharmaceutical combinations of long-acting β_2 -agonists and steroids are disclosed in two European applications, EP 416950 which discloses the combination of salmeterol and beclomethasone, and EP 416951 which discloses the combination of salmeterol and fluticasone propionate.

administration in Japan since 1986.

In Ann. Allergy 1989, 63 (3), p. 220-224 the use of a β_2 -agonist, i.e. formoterol and a steroid, i.e. budesonide seperately are mentioned. It is not disclosed a pharmaceutical combination including both formoterol and budesonide, or the use of the two compounds in combination therapy. The use of a β_2 -agonist and a steroid separately is also mentoined in Lung (1990), 168, no. supp, p. 105-110.

Outline of the Invention

35

The present invention is based on the concept of a novel combination therapy whereby formoterol (and/or a

25

35

physiologically acceptable salt and/or solvate thereof) and budesonide are administrated simultaneously, sequentially or seperately by inhalation. This combination has not only a greater efficiency and duration of bronchodilator action but the combination 5 also has a rapid onset of action. This new feature is of utmost importance in order to establish a higher compliance for patients and it provides a rescue medicine thereby avoiding the necessity for the patient of carrying two different inhalers. This simplifies life for 10 patients considerably and makes life more comfortable and secure. The rapid onset of the long-acting β_2 -agonist gives the patient immediate confirmation that he has taken an adequate dose and thereby avoiding overdosing of both $\beta_2\text{-agonist}$ and steroid. Since the use of formoterol 15 instead of salmoterol gives a much more rapid onset the combinations according to the invention have a number of advantages compared to the combinations disclosed i EP 416950 and EP 41651. The combination according to present invention permits a twice daily dosing regime as a basic 20 treatment of asthma, particularly nocturnal asthma.

The present invention provides a medicament containing, separately, or together, (i) formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and (ii) budesonide for simultaneous, sequential or separate administration by inhalation in the treatment of respiratory disorder.

The invention also provides a pharmaceutical composition for administration by inhalation in the treatment of respiratory disorder which composition comprises formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide.

According to another aspect of the invention there are provided pharmaceutical compositions comprising effective

5

10

30

amounts of formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide as a combined preparation for simultaneous, sequential or seperate administration by inhalation in the treatment of respiratory disorder.

The invention further provides formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide for use in combination therapy by simultaneous, sequential or seperate administration by inhalation in the treatment of respiratory disorder.

Further the invention provides the use of formoterol (and/or a physiologically acceptable salt and/or solvate 15 thereof) in the manufacture of a medicament for combination therapy where formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide are administered simultaneously, sequentially or seperately by inhalation in the treatment 20 of respiratory disorder and the use of budesonide in the manufacture of a medicament for combination therapy where formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide are administered simultaneously, sequentially or separately by inhalation in the treatment of respiratory disorder. 25

The invention additionally relates to the use of formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide in the manufacture of a medicament for combination therapy for simultaneous, sequential or seperate administration of formoterol and budesonide by inhalation in the treatment of respiratory disorder.

According to a further feature of the invention there is provided a method of treating respiratory disorder which comprises the simultaneous, sequential or separate

20

25

30

35

administration by inhalation of effective amounts of formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide.

Suitable physiologically salts of formoterol include acid addition salts derived from inorganic and organic acids, such as the hydrochloride, hydrobromide, sulphate, phosphate, maleate, fumarate, tartrate, citrate, benzoate, 4-methoxybenzoate, 2- or 410 hydroxybenzoate, 4-chlorobenzoate, p-toluenesulphonate, methanesulphonate, ascorbate, salicylate, acetate succinate, lactate, glutarate, gluconate, tricarballylate, hydroxynaphthalenecarboxylate or oleate. Formoterol is preferably used in the form of its fumarate salt and as a dihydrate.

The ratio of formoterol to budesonide used according to the invention is preferably within the range of 1:4 to 1:70. The two drugs may be administered separately in the same ratio.

The intended dose regimen is a twice daily administration, where the suitable daily dose of formoterol is in the range of 6 to 100 μg with a preferred dose of 6-48 μg and the suitable daily dose for budesonide is 50 to 4800 μg with a preferred dose of 100-1600 μg . The particular dose used will strongly depend on the patient (age, weight etc) and the severity of the disease (mild, moderate, severe asthma etc).

For administration, the combination is suitably inhaled from a nebulizer, from a pressurized metered dose inhaler or as a dry powder from a dry powder inhaler (e.g. as sold under the trade mark Turbuhaler) or from a dry powder inhaler utilizing gelatine, plastic or other capsules, cartridges or blister packs.

A diluent or carrier, generally non-toxic and chemically inert to the medicament e.g. lactose, dextran, mannitol or glucose or any additives that will give the medicament a desired taste, can be added to the powdered medicament.

5

10

Examples of the preparation of suitable dosage forms according to the invention include the following:
Formoterol fumarate dihydrate and budesonide (optionally premicronized) are mixed in the proportions given above. The agglomerated, free-flowing micronized mixture may be filled into dry powder inhaler such as sold under the trade mark Turbuhaler. When a capsule system issued, it is desirable to include a filler in the mixture.

- The micronized mixture may be suspended or dissolved in a 15 liquid propellant mixture which is kept in a container that is sealed with a metering valve and fitted into a plastic actuator. The propellants used may be chlorofluorocarbons of different chemical formulae. The 20 most frequently used chlorofluorocarbon propellants are trichloromonofluoromethane (propellant 11), dichlorodifluoromethane (propellant 12), dichlorotetrafluoroethane (propellant 114), tetrafluoroethane (propellant 134a) and 1,1-difuoroethane (propellant 152a). Low concentrations of a 25 surfactant such as sorbitan trioleate, lecithin, disodium dioctylsulphosuccinate or oleic acid may also be used to
- The invention is further illustrated by way of example with reference to the following Examples.

Example 1 - Dry Powder Inhaler (Turbuhaler)

improve the physical stability.

35	Active ingredient			<u>Per dos</u>	<u>se</u>	
•	Formoterol	(as	fumarate	dihydrate)	12	μg
•	Budesonide				200	μg

 $25 - 100 \mu l$

Propellant

The storage unit of the inhaler is filled with sufficient for at least 200 doses.

5	Active ingredient	<u>Per dose</u>
J	Formoterol (as fumarate dihydrate)	24 μg
	Budesonide	200 µg
	The storage unit is filled with sufficient	for at least
	200 doses.	
10	200 2000	•
10	Active ingredient	<u>Per dose</u>
	Formoterol (as fumarate dihydrate)	12 µg
	Budesonide	100 µg
	The storage unit is filled with sufficient	for at least
15	200 doses.	
10	200 doses.	
	Example 2 - Metered dose inhaler	•
	Indinote 2	
	Active ingredient	Per dose
20	Formoterol (as fumarate dihydrate)	12 µ g
20	Budesonide	200 μ g
	Stabilizer	0.1 - 0.7 mg
	Propellant	25 - 100 μl
	Floberiumė	
25		
2.7	Active ingredient	Per dose
	Formoterol (as fumarate dihydrate)	24 µg
	Budesonide	200 µ g
	Stabilizer	0.1 - 0.7 mg
20	Propellant	25 - 100 μl
30	Propertanc	•
	Active ingredient	Per dose
	Formoterol (as fumarate dihydrate)	<u></u> 12 μg
a =	Budesonide	200 μg
35		0.1 - 0.7 mg
	Stabilizer	100 13

9

Example 3 - Metered dose dry powder formulation

	Active ingredient					Per do	
	Formoterol (as fumarate dihyd	irate)				12	μg
5	Budesonide					200	μg
	Lactose	u	o to	5,	12.5	or 25	mg
,	Active ingredient					Per do	se
10	Formoterol (as fumarate dihyd	irate)				24	μg
	Buđesoniđe					200	μg
	Lactose	u	o to	5,	12.5	or 25	mg
							-
15	Active ingredient					Per do	se
	Formoterol (as fumarate dihyo	lrate)				12	μg
	Budesonide					100	μg
	Lactose	u	p to	5,	12.5	or 25	mg

20

10

CLAIMS

1. A medicament containing, seperately or together, (i) formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and (ii) budesonide for simultaneous, sequential or seperate administration by inhalation in the treatment of respiratory disorder.

5

25

35

- A pharmaceutical composition for administration by inhalation in the treatment of respiratory disorder which composition comprises formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide.
- 3. A pharmaceutical composition comprising effective amounts of formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide as a combined preparation for simultaneous, sequential or separate administration by inhalation in the treatment of respiratory disorder.
 - 4. Formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide for use in combination therapy by simultaneous, sequential or separate administration by inhalation in the treatment of respiratory disorder.
- 5. The use of formoterol (and/or a physiologically acceptable salt thereof) in the manufacture of a
 medicament for combination therapy where formoterol
 (and/or physiologically acceptable salt and/or solvate thereof) and budesonide are administered simultaneously, sequentially or separately by inhalation in the treatment of respiratory disorder.
 - 6. The use of budesonide in the manufacture of a medicament for combination therapy where formoterol

(and/or a physiologically acceptable salt and/or solvate thereof) and budesonide are administered simultaneously, sequentially or separately by inhalation in the treatment of respiratory disorder.

11

5

10

7. The use of formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide in the manufacture of a medicament for combination therapy for simultaneous, sequential or separate administration of formoterol and budesonide by inhalation in the treatment of respiratory disorder.

			international rippinearon i to		
I. CLASS	HICATION OF SUBJ	ECT MATTER (if several classification	symbols apply, indicate all) ⁶		
		t Classification (IPC) or to both National			
[Int.C]	I. 5 A61K31/5	7; //(A61K31/57	7,31:165)		
Ì		•			
II. FIELD	S SEARCHED				
	······································	Minimum Docu	mentation Searched ⁷		
Classifica	ation System		Classification Symbols		
Int.C1	1. 5	A61K			
			er than Minimum Documentation s are Included in the Fields Searched ⁸		
		to the Extent that such Document	2 ALE INCIDUES IN the Lienz Dealchen		
m. Docu	IMENTS CONSIDERE	D TO BE RELEVANT ⁹			
Category o	Citation of Do	ocument, 11 with indication, where approp	riate, of the relevant passages 12	Relevant to Claim No. ¹³	
		,			
Х	ANNALS (OF ALLERGY		1-7	
l		, no. 3, September 198	9,		
	pages 22				
		MARSAC ET AL. 'Inhale			
		and inhaled steroids	in the		
		it of asthmal			
		n the application e 221, column 2; table	1	•	
		221, column 2, table 221, column 3, line			
		223; summary *	11 17/10 10		
	P-3- "				
Х	LUNG (US	SA)		1-7	
l	vol. 168	3, no. SUPP, 1990, NEW	YORK		
	pages 10				
		DMYR 'The current pla			
		nists in the manageme	nt of asthma'		
		the application	•		
	see abst	ract 			
°.Specia	il categories of cited doc	uments : 10	"T" later document published after the interna	tional filing date	
"A" do	cument defining the gen-	eral state of the art which is not	or priority date and not in conflict with the cited to understand the principle or theory	e application but	
	nsidered to be of particu	lar relevance shed on or after the international	invention		
fili	ing date		"X" document of particular relevance; the claim cannot be considered novel or cannot be o		
· "L" doc whi	cument which may throw ich is cited to establish t	doubts on priority claim(s) or the publication date of another	invoive an inventive step		
cita	ation or other special rea	son (as specified)	"Y" document of particular relevance; the clair cannot be considered to involve an inventi	ve step when the	
	cument referring to an o her means	ral disclosure, use, exhibition or	document is combined with one or more o ments, such combination being obvious to		
P" doc	"P" document published prior to the international filing date but in the art.				
·	······································		"&" document member of the same patent fam		
IV. CERTI	FICATION				
Date of the	Actual Completion of th	e International Search	Date of Mailing of this International Sear	ch Report	
	O7 APR	IL 1993	12814146		
			100/110		
Internationa	I Searching Authority		Signature of Authorized Officer	i	
	EUROPEA	N PATENT OFFICE	LEHERTE C.F.M.	i	

Ferm PCT/ISA/210 (second sheet) (January 1985)

2



SECOND CORRECTED VERSION **

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT) **WO 93/11773** (51) International Patent Classification 5: (11) International Publication Number: A1 A61K 31/57 // (A61K 31/57 (43) International Publication Date: 24 June 1993 (24.06.93) A61K 31:165) PCT/EP92/02826 (81) Designated States: AT, AU, BB, BG, BR, CA, CH, CS, (21) International Application Number: DE, DK, ES, FI, GB, HU, JP, KP, KR, LK, LU, MG, MN, MW, NL, NO, NZ, PL, RO, RU, SD, SE, UA, (22) International Filing Date: 7 December 1992 (07.12.92) European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ. CF. CG. CI, CM, GA, GN, ML, MR, SN, TD, TG). (30) Priority data: 91311761.0 18 December 1991 (18.12.91) (34) Countries for which the regional Published or international application With international search report. SE et al. was filed: Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments. '(71) Applicant: AKTIEBOLAGET ASTRA [SE/SE]; S-151 85 Södertälje (SE). (72) Inventors: CARLING, Christer, Carl, Gustav; Backvägen 8, S-240 10 Dalby (SE). TROFAST, Jan, William; Vapenkroken 34, S-226 47 Lund (SE). (74) Agents: HJERTMAN, Ivan et al.; AB Astra, Patent Department, S-151 85 Södertälje (SE).

(54) Title: NEW COMBINATION OF FORMOTEROL AND BUDESONIDE

(57) Abstract

Effective amounts of formoterol (and/or a physiologically acceptable salt and/or solvate thereof) and budesonide are used in combination for simultaneous, sequential or separate administration by inhalation in the treatment of respiratory disorder.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinca	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU -	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic	SD	Sudan
CG	Congo		of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SI	Slovenia
Cl	Côte d'Ivoire	KZ	Kazakhstan	SK	Slovakia
CM	Cameroon	LI	Licchtenstein	SN	Senegal
CN	China	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
cz	Czech Republic	LV	Latvia	TJ	Tajikistan
DE	Germany	MC	Monaco	TT	Trinidad and Tobago
DK	Denmark	MD	Republic of Moldova	UA	Ukraine
ES	Spain	MG	Madagascar	US	United States of America
FI	Finland	. ML	Mali	UZ	Uzbekistan
FR	France	MN	Mongolia	VN	Vict Nam
GA	Gabon				